

## 兔抗 CBL (phospho-Tyr700) 多克隆抗体

中文名称: 兔抗 CBL (phospho-Tyr700) 多克隆抗体

英文名称: Anti-CBL (phospho-Tyr700) rabbit polyclonal antibody

别名: CBL2; NSLL; C-CBL; RNF55; FRA11B

相关类别: 一抗

储存: 冷冻 (-20℃) 避光

宿主: Rabbit

抗原: CBL (phospho-Tyr700)

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

### 技术规格

<b>Background:</b>	Participates in signal transduction in hematopoietic cells. Adapter protein that functions as a negative regulator of many signaling pathways that start from receptors at the cell surface. Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including PDGFA, EGF and CSF1, and terminates signaling.
<b>Applications:</b>	WB, IHC, IF
<b>Name of antibody:</b>	CBL (phospho-Tyr700)
<b>Immunogen:</b>	Synthetic peptide of human CBL (phospho-Tyr700)
<b>Full name:</b>	Cbl proto-oncogene, E3 ubiquitin protein ligase (phospho-Tyr700)

	o-Tyr700)
<b>Synonyms :</b>	CBL2; NSLL; C-CBL; RNF55; FRA11B
<b>SwissProt:</b>	P22681
<b>IHC positive control:</b>	Human breast carcinoma
<b>IHC Recommend dilution:</b>	50-100
<b>WB Predicted band size:</b>	120 kDa
<b>WB Positive control:</b>	HT29 cells treated with UV
<b>WB Recommended dilution:</b>	500-1000
<b>IF Positive control:</b>	Hela cells
<b>IF Recommended dilution</b>	100-200



