

兔抗 HDAC2 (Phospho-Ser394)多克隆抗体

中文名称： 兔抗 HDAC2 (Phospho-Ser394)多克隆抗体

英文名称： Anti-HDAC2 (Phospho-Ser394) rabbit polyclonal antibody

别 名： HD2; RPD3; YAF1

相关类别： 一抗

储 存： 冷冻 (-20°C) 避光

宿 主： Rabbit

抗 原： HDAC2 (Phospho-Ser394)

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes
Applications:	WB, IHC, IF
Name of antibody:	HDAC2 (Phospho-Ser394)
Immunogen:	Synthetic peptide of human HDAC2 (Phospho-Ser394)
Full name:	histone deacetylase 2 (Phospho-Ser394)
Synonyms :	HD2; RPD3; YAF1
SwissProt:	Q92769
IHC positive control:	Human breast carcinoma

IHC Recommend dilution:	50-100
WB Predicted band size:	55 kDa
WB Positive control:	293 cells untreated or treated with EGF
WB Recommended dilution:	500-1000
IF Positive control:	HeLa cells
IF Recommended dilution	100-200



