

兔抗 CLIP2 多克隆抗体

中文名称：兔抗 CLIP2 多克隆抗体

英文名称：Anti-CLIP2 rabbit polyclonal antibody

别名：CAP-Gly domain containing linker protein 2 CLIP; CYLN2; WSCR3; WSCR4; WBSCR3; WBSCR4; CLIP-115

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CLIP2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	The protein encoded by this gene belongs to the family of cytoplasmic linker proteins, which have been proposed to mediate the interaction between specific membranous organelles and microtubules. This protein was found to associate with both microtubules and an organelle called the dendritic lamellar body. This gene is hemizygotously deleted in Williams syndrome, a multisystem developmental disorder caused by the deletion of contiguous genes at 7q11.23. Alternative splicing of this gene generates 2 transcript variants.
Applications:	ELISA, WB, IHC

Name of antibody:	CLIP2
Immunogen:	Synthetic peptide of human CLIP2
Full name:	CAP-Gly domain containing linker protein 2
Synonyms:	CLIP; CYLN2; WSCR3; WSCR4; WBSCR3; WBSCR4; CLIP-115
SwissProt:	Q9UDT6
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer and Human ovarian cancer
IHC Recommend dilution:	20-100
WB Predicted band size:	116 kDa
WB Positive control:	LOVO and NIH/3T3 cell lysates
WB Recommended dilution:	200-1000



