

兔抗 CYSLTR2 多克隆抗体

中文名称：兔抗 CYSLTR2 多克隆抗体

英文名称：Anti-CYSLTR2 rabbit polyclonal antibody

别名：cysteinyl leukotriene receptor 2 HG57; CYSLT2; GPCR21; HPN321; CYSLT2R; KPG_011; hGPCR21; PSEC0146

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CYSLTR2

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	The cysteinyl leukotrienes LTC ₄ , LTD ₄ , and LTE ₄ are important mediators of human bronchial asthma. Pharmacologic studies have determined that cysteinyl leukotrienes activate at least 2 receptors, the protein encoded by this gene and CYSLTR1. This encoded receptor is a member of the superfamily of G protein-coupled receptors. It seems to play a major role in endocrine and cardiovascular systems.
Applications:	ELISA, WB, IHC
Name of antibody:	CYSLTR2
Immunogen:	Synthetic peptide of human CYSLTR2

Full name:	cysteinyl leukotriene receptor 2
Synonyms:	HG57; CYSLT2; GPCR21; HPN321; CYSLT2R; KPG_011; hGPCR21; PSEC0146
SwissProt:	Q9NS75
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human gastric cancer and Human liver cancer
IHC Recommend dilution:	20-100
WB Predicted band size:	40 kDa
WB Positive control:	Mouse adrenal gland tissue lysate
WB Recommended dilution:	200-1000



