

全国订货电话 4008-723-722

兔抗 DCLK2 多克隆抗体

- 中文名称: 兔抗 DCLK2 多克隆抗体
- 英文名称: Anti-DCLK2 rabbit polyclonal antibody
- 别名: CL2; DCK2; CLIK2; DCDC3; CLICK2; DCDC3B; DCAMKL2; CLICK-II
- 相关类别: 一抗
- 储存: 冷冻(-20℃) 避光
- 抗原: DCLK2
- 宿 主: Rabbit
- 反应种属: Human Mouse
- 标记物: Unconjugate
- 克隆类型: Unconjugate

技术规格

Background:	This gene encodes a member of the protein kinase super family and the doublecortin family. The protein encoded by this gene contains two N-terminal doublecortin domai ns, which bind microtubules and regulate microtubule pol ymerization, a C-terminal serine/threonine protein kinase domain, which shows substantial homology to Ca2+/calm odulin-dependent protein kinase, and a serine/proline-rich domain in between the doublecortin and the protein kina se domains, which mediates multiple protein-protein inter actions. The microtubule-polymerizing activity of the enco
	se domains, which mediates multiple protein-protein inter actions. The microtubule-polymerizing activity of the enco
	ded protein is independent of its protein kinase activity. Mouse studies show that the DCX gene, another family m ember, and this gene share function in the establishment



全国订货电话 4008-723-722

	of hippocampal organization and that their absence result s in a severe epileptic phenotype and lethality, as describ ed in human patients with lissencephaly. Multiple alternati vely spliced transcript variants have been identified.
Applications:	WB
Name of antibody:	DCLK2
Immunogen:	Synthesized peptide derived from internal of human DCLK 2.
Full name:	doublecortin-like kinase 2
Synonyms :	CL2; DCK2; CLIK2; DCDC3; CLICK2; DCDC3B; DCAMKL2; CL ICK-II
SwissProt:	Q8N568
WB Predicted band size:	84 kDa
WB Positive control:	HepG2 cells lysate
WB Recommended dilution:	500-3000

