

BMPR2 抗原(重组蛋白)

中文名称: BMPR2 抗原 (重组蛋白)

英文名称: BMPR2 Antigen (Recombinant Protein)

别 名: bone morphogenetic protein receptor, type II (serine/threonine kinase); BMR2; PPH1; BMPR3; BRK-3; T-ALK; BMPR-II

储存: 冷冻(-20℃)

相关类别: 抗原

概 述:

Fusion protein corresponding to a region derived from C terminal 250 amino acids of human BMPR2

技术规格:

| Full name: | bone morphogenetic protein receptor, type II (serine/threonine k inase) |
|--------------------|---|
| Synonyms: | BMR2; PPH1; BMPR3; BRK-3; T-ALK; BMPR-II |
| Swissprot: | Q13873 |
| Gene Accession: | BC052985 |
| Purity: | >85%, as determined by Coomassie blue stained SDS-PAGE |
| Expression system: | Escherichia coli |
| Tags: | His tag C-Terminus, GST tag N-Terminus |
| Background: | This gene encodes a member of the bone morphogenetic protei n (BMP) receptor family of transmembrane serine/threonine kina ses. The ligands of this receptor are BMPs, which are members of the TGF-beta superfamily. BMPs are involved in endochondral bone formation and embryogenesis. These proteins transduce th eir signals through the formation of heteromeric complexes of t wo different types of serine (threonine) kinase receptors: type I r |



全国订货电话 4008-723-722

eceptors of about 50-55 kD and type II receptors of about 70-8 0 kD. Type II receptors bind ligands in the absence of type I re ceptors, but they require their respective type I receptors for sig naling, whereas type I receptors require their respective type II r eceptors for ligand binding. Mutations in this gene have been a ssociated with primary pulmonary hypertension, both familial an d fenfluramine-associated, and with pulmonary venoocclusive dis ease.