

## 兔抗 FGR 多克隆抗体

中文名称：兔抗 FGR 多克隆抗体

英文名称：Anti-FGR rabbit polyclonal antibody

别名：FGR proto-oncogene, Src family tyrosine kinase; SRC2; c-fgr; c-src2; p55-Fgr; p58-Fgr; p55c-fgr; p58c-fgr

抗原：FGR

相关类别：一抗

储存：冷冻 (-20℃)

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

#### Background:

This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein contains N-terminal sites for myristylation and palmitylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to plasma membrane ruffles, and functions as a negative regulator of cell migration and adhesion triggered by the beta-2 integrin signal transduction pathway. Infection with Epstein-Barr virus results in the overexpression of this gene. Multiple alternatively spliced

	variants, encoding the same protein, have been identified.
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	FGR
<b>Immunogen:</b>	Synthetic peptide of human FGR
<b>Full name:</b>	FGR proto-oncogene, Src family tyrosine kinase
<b>Synonyms:</b>	SRC2; c-fgr; c-src2; p55-Fgr; p58-Fgr; p55c-fgr; p58c-fgr
<b>SwissProt:</b>	P09769
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human liver cancer
<b>IHC Recommend dilution:</b>	30-150

