

兔抗 PLAUR 多克隆抗体

中文名称：兔抗 PLAUR 多克隆抗体

英文名称：Anti-PLAUR rabbit polyclonal antibody

别名：CD87; UPAR; URKR; U-PAR

抗原：PLAUR

储存：冷冻（-20℃）

宿主：Rabbit

相关类别：一抗

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes the receptor for urokinase plasminogen activator and, given its role in localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the extracellular matrix. It binds both the proprotein and mature forms of urokinase plasminogen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encodi

	ng different isoforms. The proprotein experiences several post-translational cleavage reactions that have not yet been fully defined.
Applications:	ELISA, WB, IHC
Name of antibody:	PLAUR
Immunogen:	Fusion protein of human PLAUR
Full name:	Plasminogen activator, urokinase receptor
Synonyms :	CD87; UPAR; URKR; U-PAR
SwissProt:	Q03405
ELISA Recommended dilution:	2000-5000
IHC positive control:	Human lung cancer and Human colon cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	37 kDa
WB Positive control:	A549 cells
WB Recommended dilution:	500-2000



