

兔抗 TNK2 多克隆抗体

中文名称：兔抗 TNK2 多克隆抗体

英文名称： Anti-TNK2 rabbit polyclonal antibody

别名： ACK, ACK1, ACK-1, p21cdc42Hs

抗 原： TNK2

储 存： 冷冻（-20℃）

宿 主： Rabbit

反应种属： Human, Mouse, Rat

相关类别： 一抗

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

This gene encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length

	nature of only two transcript variants has been determined.
Applications:	ELISA, WB, IHC
Name of antibody:	TNK2
Immunogen:	Synthetic peptide of human TNK2
Full name:	tyrosine kinase, non-receptor, 2
Synonyms :	ACK, ACK1, ACK-1, p21cdc42Hs
SwissProt:	Q07912
ELISA Recommended dilution:	2000-10000
IHC positive control:	Human breast cancer and Human liver cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	119 kDa
WB Positive control:	231 cells
WB Recommended dilution:	1000-5000



